arXiv:1610.05827v1 [math.DS] 18 Oct 2016

A MULTIFRACTAL ANALYSIS FOR CUSPIDAL WINDINGS ON HYPERBOLIC SURFACES

JOHANNES JAERISCH, MARC KESSEBÖHMER, AND SARA MUNDAY

ABSTRACT. In this paper we investigate the multifractal decomposition of the limit set of a finitely generated, free Fuchsian group with respect to the mean cusp winding number. We will completely determine its multifractal spectrum by means of a certain free energy function and show that the Hausdorff dimension of sets consisting of limit points with the same scaling exponent coincides with the Legendre transform of this free energy function.

1. INTRODUCTION AND STATEMENT OF RESULTS

In this paper we consider a multifractal analysis of cusp windings of the geodesic flow on $\mathbb{H}/G$, for a finitely generated, free, non-elementary Fuchsian group G with parabolic elements acting on the upper half-space model $(\mathbb{H},d)$ of 2-dimensional hyperbolic space. Recall that to each ξ in the radial limit set $L_r(G)$ of G one can associate an infinite word whose letters come from the symmetric set G_0 of generators of G. Note that G can be written as a free product $G = H * \Gamma$, where $H = \langle h_1\rangle * \ldots * \langle h_u\rangle$ denotes the free product of finitely many elementary hyperbolic groups, and $\Gamma = \langle \gamma_1\rangle * \ldots * \langle \gamma_v\rangle$ denotes the free product of finitely many parabolic subgroups of G such that $\langle \gamma_i\rangle$ is the parabolic subgroup of G associated with the parabolic fixed point p_i (see also [KS04]). We will always assume that $v \geq 1$; furthermore, the fact that G is non-elementary implies that $u+v > 1$. Clearly, $\langle \gamma_i\rangle \cong \mathbb{Z}$ and $\gamma_i^{\pm 1}(p_i) = p_i$, for all $i = 1,\ldots,v$. It is well known [BJ97] that the Poincaré exponent $\delta = \delta_G$ of G coincides with the Hausdorff dimension $\dim_{\mathrm{H}}(L_r(G))$ of the radial limit set of G.

There is a natural coding of the limit set by infinite sequences over the set of generators. That is, with F referring to the Dirichlet fundamental domain of G containing $i \in \mathbb{H}$, the images of F under G tessellate $\mathbb{H}$ and each side of each of the tiles is uniquely labeled by an element of G_0. The hyperbolic ray s_ξ from i towards $\xi \in L_r(G)$ must traverse infinitely many of these tiles, and the infinite word expansion associated with ξ is then obtained by progressively recording, starting at i, the generators of the sides at which s_ξ exits the tiles. In this way we derive an infinite word $\omega = (\omega_1,\omega_2,\ldots) \in G_0^{\mathbb{N}}$, which is necessarily reduced, where reduced means that $\omega_n\omega_{n+1} \neq \mathrm{id}$, $n \in \mathbb{N}$. We then form a sequence of blocks $(B_n)_{n\in\mathbb{N}}$ in this word as follows. Each hyperbolic generator that appears in the word is given block length 1. Further, if there is a block in which the same parabolic generator appears n times and if there is no larger block of this parabolic generator containing that block, then this block is of length n. By construction, such a block of length n corresponds to the event that the projection of s_ξ onto $\mathbb{H}/G$ spirals precisely $n-1$ times around a cusp of $\mathbb{H}/G$. This

Date: 20th October 2016.

2000 *Mathematics Subject Classification.* 11K50 primary; 37A45 11J06, 28A80 secondary.

Key words and phrases. Kleinian groups, multifractal formalism, cuspidal windings.

JJ was supported by the JSPS KAKENHI 90741869 and by the Research Fellowship of the German Research Foundation (DFG-grant Ja 2145/1-1). MK was supported by the German Research Foundation (DFG) grant *Renewal Theory and Statistics of Rare Events in Infinite Ergodic Theory* (Geschäftszeichen Ke 1440/2-1). JJ and MK were supported by the DFG Scientific Network *Skew product dynamics and multifractal analysis* of the German Research Foundation (DFG-grant Oe 538/3-1).

motivates our definition of the *cusp winding process* (a_k) by setting a_k to be equal to the length $|B_k|$ of the k-th block B_k minus one.

Our main aim is to investigate the fluctuation of a certain asymptotic exponential scaling associated to this process, thereby extending results in [JK10, JK11] for $\mathrm{PSL}_2(\mathbb{Z})$. For $x \in L_r(G)$ let $B_i = B_i(x)$ denote the block of length $|B_i|$ and, for later use, set $a_i := a_i(x) := |B_i(x)| - 1$. We say that the *mean cusp-winding number* of $x \in L_r(G)$ is given by

$$\lim_{n\to\infty} \frac{2\sum_{k=1}^{n} \log^+(|B_k|-1)}{d(B_1\cdots B_n(i), i)},$$

whenever the limit exists. Here, $\log^+(a) := \begin{cases} \log(a), & \text{when } a>0; \\ 0, & \text{when } a=0. \end{cases}$. The fluctuation of this quantity is captured in the level sets

$$\mathscr{F}_\alpha := \left\{ x \in L_r(G) \mid \lim_{n\to\infty} \frac{2\sum_{k=1}^{n} \log^+(|B_k|-1)}{d(B_1\cdots B_n(i), i)} = \alpha \right\},$$

for a prescribed scaling constant $\alpha \in \mathbb{R}$.

Fact 1.1. *The subset of the limit set*

$$L_{\mathrm{c}} = \{x \in L_r(G) \mid a_k(x) \le 1 \text{ for all } k \in \mathbb{N}\},$$

encoding those geodesic rays whose maximal winding around any given cusp is at most one, is contained in $\mathscr{F}_0$. In particular, the set of limit points $L(\langle H_0\rangle)$ that can be coded exclusively via hyperbolic elements is contained in $\mathscr{F}_0$, i.e.,

$$L(\langle H_0\rangle) \subset L_{\mathrm{c}} \subset \mathscr{F}_0.$$

Fact 1.2. *We have that the set $\left\{x \in L_r(G) \mid \lim_{n\to\infty} n^{-1}\sum_{i=1}^{n} \log a_i(x) = \infty\right\}$ is contained in $\mathscr{F}_1$. In particular, for the Jarník set*

$$\mathscr{J} := \left\{ x \in L_r(G) \mid \lim_{n\to\infty} a_n(x) = \infty \text{ and } \lim_{n\to\infty} \frac{\log a_n(x)}{d(B_1\cdots B_n(i), i)} = 0 \right\}$$

considered in [Mun12] *we have $\mathscr{J} \subset \mathscr{F}_1$ and $\dim_{\mathrm{H}}(\mathscr{J}) = 1/2$.*

Fact 1.3. *For every $x \in \mathscr{F}_1$ we have that the sequence $(a_n(x))_{n\in\mathbb{N}}$ is unbounded .*

Fact 1.4. *We have $\mathscr{F}_\alpha \neq \varnothing$ if and only if $\alpha \in [0,1]$.*

Since these level sets are generally Lebesgue null sets, the Hausdorff dimension $\dim_{\mathrm{H}}(\mathscr{F}_\alpha)$ is an appropriate quantity to measure the size of the sets $\mathscr{F}_\alpha$. In this paper we will give a complete analysis of the *cusp-winding spectrum*

$$f(\alpha) := \dim_{\mathrm{H}}(\mathscr{F}_\alpha), \qquad \alpha \in \mathbb{R}.$$

Using the Thermodynamic Formalism we will be able to express the function f on $[0,1]$ implicitly in terms of the *cusp-winding pressure function*

$$P(t,\beta) := \lim_{n\to\infty} \frac{1}{n} \log \sum_{B_1\cdots B_n \in \mathscr{G}_n} \mathrm{e}^{-td(B_1\cdots B_n(i),i) - 2\beta\sum_{i=1}^{n}\log^+(|B_i|-1)}, \quad t,\beta \in \mathbb{R},$$

where $\mathscr{G}_n := \{B_1(x)\cdots B_n(x) \in G \mid \ x \in L_r(G)\}$. The sequence $(Z_n)_n$ with

$$Z_n := \log \sum_{B_1\cdots B_n \in \mathscr{G}_n} \mathrm{e}^{-td(B_1\cdots B_n(i),i) - 2\beta\sum_{i=1}^{n}\log^+(|B_i|-1)}, \ n \in \mathbb{N},$$

is almost sub-additive in the sense that $Z_{n+m} \le Z_n + Z_m + 2|t|C_0$ for a certain constant $C_0 > 0$ which will be defined in Section 2.2. In fact, this estimate boils down to the inequality $d(B_1\cdots B_{n+m}(i), i) \ge d(B_1\cdots B_n(i), i) + d(B_1\cdots B_m(i), i) - 2C_0$, which follows from (2.1)



FIGURE 1.1. The *cusp-winding free energy function* $\beta \mapsto t(\beta)$ and the scaling spectrum $\alpha \mapsto f(\alpha)$.

below. This is a consequence of the triangle inequality and the definition of the blocks B_i. Hence the limit of (Z_n/n) exists and is equal to

$$\lim_{n\to\infty} \frac{Z_n}{n} = \inf_{k\in\mathbb{N}} \left(\frac{Z_k}{k} + \frac{2|t|C_0}{k} \right). \tag{1.1}$$

We shall see in Lemma 2.4 that P coincides with the dynamically-defined topological pressure function given in (2.4). By Proposition 2.5 we have that for every $\beta \in \mathbb{R}$ there exists a unique number $t = t(\beta)$, such that $P(t(\beta), \beta) = 0$. We denote by $\beta \mapsto t(\beta)$ the *cusp-winding free energy function* (see Fig. 1.1). For any real convex function g we let $\widehat{g}\colon \mathbb{R} \to \mathbb{R} \cup \{\infty\}$ denote the *Legendre transform of* g, which is defined to be

$$\widehat{g}(p) := \sup\{cp - g(c)\colon c \in \mathbb{R}\}, \text{ for all } p \in \mathbb{R}.$$

Further, let $\delta = \delta(G)$ denote the exponent of convergence of the Poincaré series of G. Now we are in the position to state our main theorem.

Theorem 1.5 (Multifractal cusp-winding analysis)**.** *For each free, finitely generated non-elementary Fuchsian group G with parabolic elements the following statements hold. The Hausdorff dimension for the cusp winding spectrum is given by*

$$f(\alpha) = \max\left\{-\widehat{t}(-\alpha), 0\right\}, \qquad \alpha \in \mathbb{R}.$$

The function $f\big|_{[0,1]}$ is strictly concave, continuous, real-analytic on $(0,1)$ and its maximal value is equal to $\delta(G)$ (cf. Fig 1.1). For the boundary points we have

$$f(0) = \delta_c := \dim_{\mathrm{H}}(L_c),\ f(1) = \frac{1}{2},\ \textit{and}\ \lim_{\alpha\searrow 0} f'(\alpha) = -\lim_{\alpha\nearrow 1} f'(\alpha) = +\infty.$$

Further, the set of points for which the mean cusp winding number does not exist has full Hausdorff dimension $\delta(G)$.

2. ERGODIC THEORY FOR FUCHSIAN GROUPS

In this section we study the ergodic theory for Fuchsian groups with parabolic elements. In Section 2.1 we describe the action of G on $\partial\mathbb{H}$ by a canonical Markov map which is referred to as the Bowen–Series map. In Section 2.2 we give the crucial geometric estimates which underly the whole multifractal analysis. In Section 2.3 we set up the induced dynamics of the Bowen–Series map on the complement of some neighbourhood of the parabolic fixed points of G. For the induced system we can later use Gibbs theory to prove the multifractal formalism.

2.1. **The canonical Markov map.** As already mentioned in the introduction, throughout we exclusively consider a finitely generated, free Fuchsian group G. Recall that G can be written as a free product $G = H * \Gamma$, where $H = \langle h_1\rangle * \ldots * \langle h_u\rangle$ denotes the free product of finitely many elementary, hyperbolic groups, and $\Gamma = \langle \gamma_1\rangle * \ldots * \langle \gamma_v\rangle$ denotes the free product of finitely many parabolic subgroups $\langle \gamma_i\rangle$ of G with the parabolic fixed point $p_i = \gamma_i^{\pm}(p_i)$, $i = 1,\ldots,v$. Since G is finitely generated, G admits the choice of a Poincaré polyhedron F with a finite set $\mathscr{F}$ of faces. Let us now first recall from [SS05] the construction of the relevant coding map T associated with G, which maps the radial limit set $L_r(G)$ into itself. This construction parallels the construction of the well-known Bowen–Series map (cf. [BS79], [Sta04]). For $\xi,\eta \in L_r(G)$, let $\gamma_{\xi,\eta} : \mathbb{R} \to \mathbb{H}$ denote the directed geodesic from η to ξ such that $\gamma_{\xi,\eta}$ intersects the closure $\overline{F}$ of F in $\mathbb{H}$, and normalised such that $\gamma_{\xi,\eta}(0)$ is the summit of $\gamma_{\xi,\eta}$. The exit time $e_{\xi,\eta}$ is defined by

$$e_{\xi,\eta} \coloneqq \sup\left\{s \mid \gamma_{\xi,\eta}(s) \in \overline{F}\right\}.$$

Since $\xi,\eta \in L_r(G)$, we clearly have that $|e_{\xi,\eta}| < \infty$. By Poincaré's Polyhedron Theorem (cf. [EP94]), we have that the set $\mathscr{F}$ carries an involution $\mathscr{F} \to \mathscr{F}$, given by $s \mapsto s'$ and $s'' = s$. In particular, for each $s \in \mathscr{F}$ there is a unique *face-pairing transformation* $g_s \in G$ such that $g_s(\overline{F}) \cap \overline{F} = s'$. We then let

$$\mathscr{L}_r(G) \coloneqq \{(\xi,\eta) \mid \xi,\eta \in L_r(G), \xi \neq \eta \text{ and } \exists t \in \mathbb{R} \,:\, \gamma_{\xi,\eta}(t) \in \overline{F}\},$$

and define the map $S : \mathscr{L}_r(G) \to \mathscr{L}_r(G)$, for all $(\xi,\eta) \in \mathscr{L}_r(G)$ such that $\gamma_{\xi,\eta}(e_{\xi,\eta}) \in s$, for some $s \in \mathscr{F}$, by

$$S(\xi,\eta) \coloneqq (g_s(\xi), g_s(\eta)).$$

In order to show that the map S admits a Markov partition, we introduce the following collection of subsets of the boundary $\partial\mathbb{H}$ of $\mathbb{H}$. For $s \in \mathscr{F}$, let A_s refer to the open hyperbolic half-space for which $F \subset \mathbb{H} \setminus A_s$ and $s \subset \partial A_s$. We then define the projection b_s of the side s to $\partial\mathbb{H}$ by

$$b_s \coloneqq \operatorname{Int}\left(\operatorname{Cl}_{\overline{\mathbb{H}}}(A_s) \cap \partial\mathbb{H}\right).$$

Clearly, $b_s \cap b_t = \varnothing$, for all $s,t \in \mathscr{F}, s \neq t$. Hence, by convexity of F, we have $\gamma_{\xi,\eta}(e_{\xi,\eta}) \in s$ if and only if $\xi \in b_s$. In other words, $S(\xi,\eta) = (g_s\xi, g_s\eta)$ for all $\xi \in b_s$. This immediately gives that the projection map $p_1 : (\xi,\eta) \mapsto \xi$ onto the first coordinate of $\mathscr{L}_r(G)$ leads to a canonical factor T of S, that is, we obtain the map

$$T : L_r(G) \to L_r(G), \text{ given by } T|_{b_s \cap L_r(G)} \coloneqq g_s.$$

Clearly, T satisfies $p_1 \circ S = T \circ p_1$. Since $T(b_s) = g_s(b_s) = \operatorname{Int}(\partial\mathbb{H} \setminus b_{s'})$, it follows that T is a non-invertible Markov map with respect to the partition $\{b_s \cap L_r(G) \mid s \in \mathscr{F}\}$. For this so-obtained expansive map T we then have the following result.

Proposition 2.1 ([SS05, Proposition 2, Proposition 3])**.** *The map T is a topologically mixing Markov map with respect to the partition generated by $\{b_s \cap L_r(G) \mid s \in \mathscr{F}\}$. Moreover, the map S is the natural extension of T.*

2.2. **Horocircles and basic estimates.** Recall that for $x \in L_r(G)$ we let $B_i(x)$ denote the i-th block of the infinite word expansion of the geodesic ray from i towards x and $a_i(x) = |B_i(x)| - 1$. In order to obtain estimates for $d(B_1 \cdots B_n(i), i)$, we introduce the following horocircles. For each parabolic generator $\gamma_i \in \Gamma_0$ with fixed point p_i, we define the *horocircle H_i of Euclidean height* $1/2$ which is given as follows: there exists a unique $\Delta \in \mathrm{PSL}_2(\mathbb{R})$ such that $\Delta(p_i) = \infty$ and $\Delta^{-1}\gamma_i\Delta(z) = z+1$. Then we set $H_i \coloneqq \Delta^{-1}(\{\Im\mathfrak{m}(z) = 1/2\})$. Another way to define H_i is to require that the collar geodesic, i.e. the projection of H_i to $\mathbb{H}/G$, has hyperbolic length 2. These horocircles are pairwise disjoint. To see this, first note that without loss of generality we can assume that one of the parabolic generators of G is of the form $z \mapsto z+1$, with fixed point at infinity. The fact that G is a free group ensures that the edges of the Dirichlet fundamental domain for G do not intersect. An



FIGURE 2.1. On the left we show a typical fundamental domain with two parabolic and one hyperbolic generator. On the right we show the largest possible horocircle with Euclidean height $1/2$ corresponding to the cusp at 0 just touching another horocircle with Euclidean height $1/2$ corresponding to the cusp at infinity.

example is shown in the left-hand side of Figure 2.1. Thus, in the configuration on the right of Figure 2.1 the horocircle has maximal Euclidean height. It is therefore sufficient to verify that the horocircles are disjoint in that case. Let $\Delta \in \mathrm{PSL}_2(\mathbb{R})$ be given by $\Delta(z) := -1/(4z)$. Then the geodesic u in Figure 2.1 (right) is mapped to the vertical line through $\Delta(-1/2) = 1/2$. Similarly, the geodesic u' is mapped to the vertical line through $\Delta(1/2) = -1/2$. Hence, the parabolic generator γ corresponding to the side-pairing of u and u' satisfies $\Delta^{-1}\gamma\Delta(z) = z+1$ and the dashed horocircle through 0 is represented by the horizontal line through $\Delta(i/2) = i/2$. We have thus shown that the dashed horocircle through zero has Euclidean height $1/2$.

To estimate $d(B_1 \cdots B_n(i), i)$, we will partition the directed geodesic segment ξ which goes from i to $B_1 \cdots B_n(i)$ and corresponds to $x \in L_r(G)$ into $(n+1)$ arcs $\xi_1, \ldots, \xi_{n+1}$ as follows. The first arc ξ_1 starts at i and the last arc ξ_{n+1} terminates at $B_1 \cdots B_n(i)$. For $k = 1, \ldots, n$, the arc ξ_k terminates at the first intersection of ξ and an image of a side of the Dirichlet fundamental domain corresponding to G centred in i, whenever $|B_k| = 1$, or the intersection of ξ with the first horocircle H_j, $j = 1, \ldots, \nu$, when leaving this horocircle. The terminating point of ξ_k coincides with the initial point of the following arc ξ_{k+1}.

If we cut off the cusps of $\mathbb{H}/G$ along the horocircles $H_1, \ldots, H_\nu$, then we obtain a compact manifold which in particular has a finite diameter $C_0 > 0$. Since ξ_{n+1} is contained in this compact set we have for all $n \in \mathbb{N}$ that,



FIGURE 2.2. The first three geodesic arcs ξ_1, ξ_2, ξ_3 connecting i and $g\gamma_k^2 h(i)$, $\gamma_k \in \Gamma_0$, $g,h \in H_0$.

$$\sum_{k=1}^{n} l(\xi_k) \leq d(B_1 \cdots B_n(i), i) = \sum_{k=1}^{n+1} l(\xi_k) \leq \sum_{k=1}^{n} l(\xi_k) + C_0, \tag{2.1}$$

where $l(\xi_k)$ refers to the hyperbolic length of the arc ξ_k (see Figure 2.2).

Before proving the facts stated in the introduction we will need the following geometric observation.



FIGURE 2.3. The longest possible parabolic excursion above $\{\mathfrak{Im}(z) = 1/2\}$ (dotted line) corresponding to two windings, i.e. $n = 4$ in Proposition 2.2.

Proposition 2.2. *For $n \geq 3$ we have that the length of the geodesic arc ξ lying above $\{\mathfrak{Im}(z) = 1/2\}$ and connecting $\pm(n-1)/2$ (cf. Figure 2.3) lies between the two constants $\log\left(3 \cdot (n-1)^2\right)$ and $\log\left(4 \cdot (n-1)^2\right)$.*

Proof. Let $0 < h < r$ and put $a := \sqrt{r^2 - h^2}$. We parametrise the circular arc of the circle with centre 0 and radius r which lies above the line $\{z \in \mathbb{H} \mid \Im\mathfrak{m}(z) = h\}$. The parametrisation is given by $\gamma : [-a, a] \to \mathbb{H}$, $\gamma(t) := \left(t, \sqrt{r^2 - t^2}\right)$. We have

$$\gamma'(t) = \left(1, -t(r^2 - t^2)^{-1/2}\right) \quad \text{and } \|\gamma'(t)\|_2 = \sqrt{1 + \frac{t^2}{r^2 - t^2}} = \frac{r}{\sqrt{r^2 - t^2}}.$$

Hence, the hyperbolic length of γ is given by

$$l(\gamma) = \int_{-a}^{a} \|\gamma'(t)\|_2 \frac{1}{\sqrt{r^2 - t^2}} dt = \frac{1}{r} \int_{-a}^{a} \frac{1}{1 - (t/r)^2} dt = \int_{-a/r}^{a/r} \frac{1}{1 - y^2} dy = \log\left(\frac{1 + a/r}{1 - a/r}\right).$$

It will be convenient to write

$$l(\gamma) = \log\left(\frac{1 + a/r}{1 - a/r}\right) = \log\left(\frac{(1 + a/r)^2}{1 - (a/r)^2}\right).$$

To verify the statement in the proposition, we must consider the case that $h := 1/2$ and $r := (n-1)/2$. Then $a = \sqrt{n^2 - 2n}/2$ and by our previous calculation we obtain

$$l(\xi) = \log\left(\frac{\left(1 + \frac{\sqrt{n^2 - 2n}}{n-1}\right)^2}{1 - \frac{n^2 - 2n}{(n-1)^2}}\right) = \log\left((n-1)^2 \left(1 + \frac{\sqrt{n^2 - 2n}}{n-1}\right)^2\right).$$

To complete the proof, observe that, for all $n \geq 3$, we have $4 \geq \left(1 + \frac{\sqrt{n^2 - 2n}}{n-1}\right)^2 \geq 3$. □

Corollary 2.3. *For any geodesic arc ξ_i corresponding to a given block B_i with $a_i \geq 2$, we have*

$$2\log(a_i) + \log 3 \leq l(\xi_i). \tag{2.2}$$

If $a_i \geq 0$ then we still have

$$2\log^+(a_i) + C_1 \leq l(\xi_i) \leq 2\log(a_i + 1) + C_2, \tag{2.3}$$

where $C_1 := \min\{\log 3, d(s,t) \mid s,t \in \mathscr{F}, g_s g_t \neq id.\}$ and $C_2 := \max\{\log 4, C_0\}$ with C_0 defined in Section 2.2.

2.3. **Inducing and the topological Markov chain with infinite state space .** Let us set

$$\mathscr{D} := \{y \in L_r(G) \mid a_1(y) = 0\}.$$

The induced transformation $T_{\mathscr{D}}$ on $\mathscr{D} = \{a_1 = 0\}$ is defined by $T_{\mathscr{D}}(\xi) := T^{\rho(\xi)}(\xi)$, where ρ denotes the return time function, given by $\rho(\xi) := \min\{n \in \mathbb{N} \mid T^n(\xi) \in \mathscr{D}\}$. Denote by $\Gamma_0 := \left\{\gamma_i, \gamma_i^{-1} \mid 1 \leq i \leq v\right\}$ the symmetric set of parabolic generators and denote by $H_0 := \left\{h_i, h_i^{-1} \mid 1 \leq i \leq u\right\}$ the symmetric set of hyperbolic generators. Recall that $G_0 = \Gamma_0 \cup H_0$. Define the induced partition

$$\alpha_{\mathscr{D}} := \bigcup_{n \in \mathbb{N}} \mathscr{D} \cap \{\rho = n\} \cap \bigvee_{k=0}^{n+1} T^{-k}\left(\{b_s \cap L_r(G) \mid s \in \mathscr{F}\}\right)$$

and the infinite alphabet

$$I := \left\{g_1 \gamma^n g_2 \mid \gamma \in \Gamma_0, g_1, g_2 \in G_0 \setminus \{\gamma^{\pm 1}\}, n \in \mathbb{N}\right\} \cup$$
$$\cup \left\{g_1 h g_2 \mid g_1 \in G_0, h \in H_0 \setminus \{g_1^{-1}\}, g_2 \in G_0 \setminus \{h^{-1}\}\right\}.$$

The incidence matrix $A \in \{0,1\}^{I\times I}$ is for $(\eta_1,\dots,\eta_n)\in I$ and $(\tau_1,\dots,\tau_m)\in I$, $n,m\geq 3$, given by $A((\eta_1,\dots,\eta_n),(\tau_1,\dots,\tau_m))=1$ if and only if $\eta_{n-1}=\tau_1$ and $\eta_n=\tau_2$. We consider the Markov shift (Σ_A,σ) where

$$\Sigma_A := \left\{ \omega=(\omega_1,\omega_2,\dots)\in I^{\mathbb{N}} \mid \forall k\in\mathbb{N}\ A(\omega_k,\omega_{k+1})=1 \right\}$$

and $\sigma\colon \Sigma_A\to\Sigma_A$, $(\sigma(\omega))_i=\omega_{i+1}$ for $\omega\in\Sigma_A$ denotes the *left shift map.*

Recall that $\mathscr{D}:=\{a_1=0\}$. The induced Bowen–Series map $(\mathscr{D},T_{\mathscr{D}})$ is conjugated to the left shift (Σ_A,σ) via π, that is, we have the following commutative diagram

$$\begin{array}{ccc} \mathscr{D} & \xrightarrow{T_{\mathscr{D}}} & \mathscr{D} \\ {\scriptstyle\pi}\downarrow & & \downarrow{\scriptstyle\pi} \\ \Sigma_A & \xrightarrow[\sigma]{} & \Sigma_A \end{array}$$

For $\omega\in\Sigma_A$ and $n\in\mathbb{N}$ let $\omega|_n:=(\omega_1,\dots,\omega_n)$ and let $[\omega|_n]:=\{\tau\in\Sigma_A \mid \tau_1=\omega_1,\dots,\tau_n=\omega_n\}$ denote the *n-cylinder* of ω. We have $\pi(\mathscr{D})=\{[\omega]\mid\omega\in I\}$.

We now aim to express the cusp-winding scaling limit in dynamical terms. For this we introduce the two *potential functions*

$$\begin{aligned} &\psi\colon \Sigma_A\to\mathbb{R}_0^-,\ (\omega_1,\omega_2,\dots)\mapsto -2\log^+(|\omega_1|-3),\\ &\varphi\colon \Sigma_A\to\mathbb{R}^-,\ (\omega_1,\omega_2,\dots)\mapsto -\log\left(\left|\left(g_1\cdots g_{|\omega_1|-1}\right)'\left(\pi^{-1}(\omega)\right)\right|\right),\\ &\qquad\qquad\qquad\qquad \text{with } \omega_1=g_1\cdots g_{|\omega_1|}\in I, \end{aligned}$$

where ψ describes the cusp-winding number, while φ describes the geometric properties of the geodesic flow. We will equip Σ_A with the metric d which is given for each $\omega,\tau\in I^{\mathbb{N}}$ by $d(\omega,\tau):=\exp(-|\omega\wedge\tau|)$, where $|\omega\wedge\tau|$ denotes the length of the longest common initial block of ω and τ. Since ψ depends only on the first symbol, we immediately see that ψ is locally Hölder continuous with respect to this metric. For the proof that φ is also locally Hölder continuous we refer to [KS04].

2.4. **Topological pressure.** First, let us fix some notation. For two sequences (a_n), (b_n) we will write $a_n\ll b_n$, if $a_n\leq Kb_n$ for some $K>0$ and all $n\in\mathbb{N}$, and if $a_n\ll b_n$ and $b_n\ll a_n$ then we write $a_n\asymp b_n$.

For $n\in\mathbb{N}$ put

$$\Sigma_A^n=\{\omega=(\omega_1,\dots,\omega_n)\in I^n \mid A(\omega_i,\omega_{i+1})=1 \text{ for all } 1\leq i\leq n-1\}.$$

The topological pressure $\mathfrak{P}(t\varphi+\beta\psi)$ of the potential $t\varphi+\beta\psi$ for $t,\beta\in\mathbb{R}$ is defined to be

$$\mathfrak{P}(t\varphi+\beta\psi) := \lim_{n\to\infty}\frac{1}{n}\log\sum_{\omega\in\Sigma_A^n}\exp\sup_{\tau\in[\omega]}\left(S_\omega(t\varphi+\beta\psi)\right)(\tau), \tag{2.4}$$

where we set $S_\omega f:=\sup_{x\in[\omega]}S_nf(x)$ with $S_nf(x):=\sum_{k=0}^{n-1}f\circ\sigma^k(x)$ and $S_0f=0$. By a standard argument involving sub-additivity the above limit always exists (although it is possibly equal to infinity).

The next lemma shows that the set $\mathscr{F}_\alpha$ can be characterised by the potentials φ and ψ and that the cusp-winding pressure $P(t,\beta)$ agrees with $\mathfrak{P}(t\varphi+\beta\psi)$ for all $t,\beta\in\mathbb{R}$. In the proof of the following lemma we make use of the fact that the topological Markov chain (Σ_A,σ) is *finitely primitive*, that is, there exists $\ell\in\mathbb{N}$ and a finite set $F\subset\Sigma_A^\ell$ such that for all $a,b\in I$ there exists $u\in\Sigma_A^\ell$ such that the word aub belongs to $\Sigma_A^{\ell+2}$, for more details, see [MU03].

Lemma 2.4. *For $\alpha \in \mathbb{R}$, $x \in \mathscr{D}$ and $\omega := \pi(x) \in \Sigma_A$ we have*

$$\lim_{n\to\infty} \frac{S_n\psi(\omega)}{S_n\varphi(\omega)} = \alpha \iff \lim_{n\to\infty} \frac{2\sum_{i=1}^n \log(a_i(x))}{d(B_1 \dots B_n(i), i)} = \alpha,$$

and for all $t, \beta \in \mathbb{R}$ we have

$$P(t,\beta) = \mathfrak{P}(t\varphi + \beta\psi).$$

Moreover, $\mathfrak{P}(t\varphi+\beta\psi) < \infty$ if and only if $t+\beta > 1/2$. Further, for each $\beta \in \mathbb{R}$ we have $\lim_{t\searrow \frac{1}{2}-\beta} P(t,\beta) = \infty$ and $\lim_{t\to\infty} P(t,\beta) = -\infty$.

Proof. Following [KS04], we find a constant $D > 0$ such that for all $x \in L_r(G)$, $\omega := \pi(x) \in \Sigma_A$ and $n \in \mathbb{N}$,

$$|d(B_1 \dots B_n(i), i) - S_n\varphi(\omega)| \le D. \tag{2.5}$$

This proves the first two assertions.

Since (Σ_A, σ) is finitely primitive, it follows from [MU03, Proposition 2.1.9] that

$$\mathfrak{P}(t\varphi+\beta\psi) < \infty \quad \text{if and only if} \quad \sum_{a\in I} e^{\sup_{x\in[a]} t\varphi(x)+\beta\psi(x)} < \infty.$$

Now, we deduce with (2.5), (2.1) and (2.3) that for all $t, \beta \in \mathbb{R}$

$$\sum_{a\in I} e^{\sup_{x\in[a]} t\varphi(x)+\beta\psi(x)} \asymp \sum_{n\ge 1} e^{-2t\log(n)-2\beta\log(n)} \asymp \sum_{n\ge 1} n^{-2(\beta+t)}. \tag{2.6}$$

which converges if and only if $t+\beta > 1/2$. This also shows $\lim_{t\searrow\frac{1}{2}-\beta} P(t,\beta) = \infty$.

Finally, by (1.1) and Corollary 2.3 (with the constant C_1 as defined in Corollary 2.3), we have for fixed $n > C_1/(2C_0)$ and $t > 0$,

$$\begin{aligned} P(t,\beta) &\le \frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} e^{-td(B_1\cdots B_n(i),i)-2\beta\sum_{i=1}^n \log^+(|B_i|-1)} + \frac{2tC_0}{n} \\ &\le \frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} e^{-t\sum_{i=1}^n \left(2\log^+(a_i)+C_1\right)-2\beta\sum_{i=1}^n \log^+(a_i)} + \frac{2tC_0}{n} \\ &\le \underbrace{t\left(\frac{2C_0}{n} - C_1\right)}_{<0} + \log\left(2\nu\sum_{k=1}^{\infty} k^{-2(t+\beta)} + 2u\right) \to -\infty \text{ for } t\to\infty. \end{aligned}$$

□

2.5. **The cusp-winding free energy.** The following proposition shows the existence of the cusp-winding free energy function and some of its basic properties.

Proposition 2.5. *For each $\beta \in \mathbb{R}$ there exists a unique number $t(\beta)$ such that*

$$P(t(\beta), \beta) = 0. \tag{2.7}$$

The cusp-winding free energy function $t : \mathbb{R} \to \mathbb{R}$ defined in this way is real-analytic and strictly convex.

Proof. We have to verify that the function $(t,\beta) \mapsto P(t,\beta)$ is real-analytic on the set $\{(t,\beta) \in \mathbb{R}^2 \mid 2(t+\beta) > 1\}$. Recall that Σ_A is a finitely primitive Markov shift. Let $(t,\beta) \in \mathbb{R}^2$ with $2(t+\beta) > 1$. By [MU03, Proposition 2.6.13] it suffices to show that $\varphi, \psi \in \mathscr{L}(\Sigma_A, \mu_{t\varphi+\beta\psi})$. Here, $\mu_{t\varphi+\beta\psi}$ denotes the unique equilibrium state of $t\varphi+\beta\psi$ for the dynamical system (Σ_A, σ), i.e., $\mu_{t\varphi+\beta\psi}$ is the unique σ-invariant Borel probability measure on Σ_A such that $h(\mu_{t\varphi+\beta\psi}) + \int t\varphi+\beta\psi \, d\mu_{t\varphi+\beta\psi} = 0$, where $h(\mu_{t\varphi+\beta\psi})$ refers to the metric entropy of the measure-theoretic dynamical system $(\Sigma_A, \sigma, \mu_{t\varphi+\beta\psi})$. It is well known that

$\mu_{t\varphi+\beta\psi}$ is a Gibbs measure with respect to $t\varphi+\beta\psi$. In particular, using the estimate in (2.6), we have that $\mu_{t\varphi+\beta\psi}([\omega_1]) \ll \exp(-2t\log|\omega_1|-2\beta\log|\omega_1|)$ for all $\omega_1 \in I$. Since $2(t+\beta)>1$, it follows that $\varphi,\psi \in \mathscr{L}(\Sigma_A,\mu_{t\varphi+\beta\psi})$. By [MU03, Theorem 2.6.12] we know that the pressure P is real-analytic on $\{(t,\beta)\in\mathbb{R}^2 \mid P(t,\beta)<\infty\}$, which by Lemma 2.4 is equal to $\{(t,\beta)\in\mathbb{R}^2 \mid 2(t+\beta)>1\}$.

Now let $\beta\in\mathbb{R}$. By Lemma 2.4 we have that $P(t,\beta)<\infty$, if and only if $t>1/2-\beta$. Also by Lemma 2.4, we have $\lim_{t\searrow\frac{1}{2}-\beta} P(t,\beta)=\infty$ and $\lim_{t\to\infty} P(t,\beta)=-\infty$. Hence, we conclude that there exists a unique $t=t(\beta)$ with $P(t(\beta),\beta)=0$.

The strict convexity of t follows from convexity and real-analyticity, unless t' is constant. That t' is not constant follows from the asymptotic behaviour of t for $\beta\to\pm\infty$ as described in the following Lemmas 2.6 and 2.7. □

Lemma 2.6. *For the right asymptotic of t we have that* $\lim_{\beta\to\infty} t(\beta)=\delta_c$.

Proof. For $s\in\mathbb{R}$ and $n\in\mathbb{N}$, set

$$p_n(s) := \frac{1}{n}\log \sum_{\substack{B_1\cdots B_n\in\mathscr{G}_n\\ |B_i|\le 2}} \mathrm{e}^{-sd(B_1\cdots B_n i,i)} \text{ and } p(s) := \lim_n p_n(s).$$

Then for all $\beta>0$, we have

$$0=\lim_{n\to\infty}\frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} \mathrm{e}^{-t(\beta)d(B_1\cdots B_n i,i)-2\beta\sum\log^+(a_i)} \ge \lim_{n\to\infty} p_n(t(\beta)) = p(t(\beta)).$$

By the formula of Bishop and Jones ([BJ97]) in follows that $t(\beta)\ge\delta_c$.

Now, let $C:=2C_0$ with C_0 as in Section 2.2, and recall that the constant ν denotes the number of cusps. Let ζ denote the Riemann zeta-function and choose $\varepsilon>0$. Then by using the triangle inequality $3(n-k)$ times corresponding to blocks of length greater than 4, we get

$$\begin{aligned}
&\frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} \mathrm{e}^{-(\delta_c+\varepsilon)d(B_1\cdots B_n i,i)-2\beta\sum\log^+(a_i)}\\
&\le \frac{1}{n}\log\sum_{k=0}^{n}\binom{n}{k}\mathrm{e}^{kp_k(\delta_c+\varepsilon)}\mathrm{e}^{3(\delta_c+\varepsilon)C(n-k)} \sum_{\substack{(B_1,\dots,B_{n-k})\in\mathscr{G}_1^{n-k}\\ |B_i|\ge 3}} \mathrm{e}^{\sum_{\ell=1}^{n-k}-(\delta_c+\varepsilon)d(B_\ell i,i)-2\beta\log^+(a_i)}\\
&\le \frac{1}{n}\log\sum_{k=0}^{n}\binom{n}{k}\mathrm{e}^{kp_k(\delta_c+\varepsilon)}\mathrm{e}^{3(\delta_c+\varepsilon)C(n-k)} \sum_{\substack{(B_1,\dots,B_{n-k})\in\mathscr{G}_1^{n-k}\\ |B_i|\ge 3}} \prod_{i=1}^{n-k} a_i^{-2\beta}\\
&\le \frac{1}{n}\log\sum_{k=0}^{n}\binom{n}{k}\mathrm{e}^{kp_k(\delta_c+\varepsilon)}\mathrm{e}^{3(\delta_c+\varepsilon)C(n-k)} \left(2\nu(\zeta(2\beta)-1)\right)^{n-k}\\
&\xrightarrow{n\to\infty} \log\left(\underbrace{\mathrm{e}^{p(\delta_c+\varepsilon)}}_{<1}+\underbrace{2\mathrm{e}^{3(\delta_c+\varepsilon)C}\nu(\zeta(2\beta)-1)}_{\xrightarrow{\beta\to\infty}0}\right)<0
\end{aligned}$$

for β large enough. Here we used the general observation that for $\lambda_k\to\lambda>0$ and $b>0$ we have

$$\lim_{n\to\infty}\left(\sum_{k=0}^{n}\binom{n}{k}\lambda_k^k b^{n-k}\right)^{1/n}=\lambda+b.$$

This shows that for all sufficiently large β we have $\delta_c\le t(\beta)\le\delta_c+\varepsilon$. Since $\varepsilon>0$ was arbitrary, it follows that $\lim_{\beta\to\infty}t(\beta)=\delta_c$. □

Lemma 2.7. *For all $\varepsilon \in (0,1/2)$ we find that for all $-\beta$ large enough we have*

$$\frac{1}{2}-\beta < t(\beta) < \frac{1}{2}-\beta+\frac{\varepsilon}{2}.$$

Proof. Combining Lemma 2.4 and Proposition 2.5 we deduce that $t(\beta) \geq 1/2-\beta$.

For the proof of the upper bound it suffices to show that for a given $\varepsilon \in (0,1/2)$ and $t_\beta := 1/2-\beta+\varepsilon/2$ we have $P(t_\beta,\beta) < 0$ for all $-\beta$ large. Recall once more that ν denotes the number of cusps and $u = \operatorname{card}(H_0)$. Fix a natural number $n > 2C_0/C_1$ with $C_0 > 0$ taken from (1.1) and C_1 taken from (2.3). By (1.1) and (2.3) combined with (2.1) we have for all β negative

$$\begin{aligned} P(t_\beta,\beta) &\leq \frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} \mathrm{e}^{-t_\beta d(B_1\cdots B_n(i),i)-2\beta\sum_{i=1}^n \log^+(|B_i|-1)} + \frac{2t_\beta C_0}{n} \\ &\leq \frac{1}{n}\log \sum_{B_1\cdots B_n\in\mathscr{G}_n} \mathrm{e}^{-t_\beta\sum_{i=1}^n\left(2\log^+(a_i)+C_1\right)-2\beta\sum_{i=1}^n\log^+(a_i)} + \frac{2t_\beta C_0}{n} \\ &\leq \underbrace{t_\beta\left(\frac{2C_0}{n}-C_1\right)}_{<0} + \frac{1}{n}\log\left(\left(2\nu\sum_{k=1}^{\infty}k^{-(1+\varepsilon)}+2u\right)^n\right) \to -\infty \text{ for } \beta\to-\infty. \end{aligned}$$

□

3. Proof of facts

Proof of Fact 1.1. The inclusion $L_c \subset \mathscr{F}_0$ follows immediately from the definition of $\mathscr{F}_0$. □

Proof of Fact 1.2. Using (2.1) and (2.3) along with the property that $\log(k+1) \leq \log^+(k)+1$, for $k \in \mathbb{N}_0$, we have

$$\begin{aligned} \liminf_{n\to\infty}\frac{2\sum_{i=1}^n\log^+(a_i(x))}{d(B_1\cdots B_n(i),i)} &\geq \liminf_{n\to\infty}\frac{\sum_{i=1}^n\log^+(a_i(x))}{\sum_{i=1}^n\log^+(a_i(x))+n(C_2+1)+C_0/2} \\ &= \liminf_{n\to\infty}\left(1+\frac{n(C_2+1)+C_0/2}{\sum_{i=1}^n\log^+(a_i(x))}\right)^{-1}. \end{aligned}$$

By assumption, the Cesàro average of $\log(a_i(x))$ tends to infinity, which implies that

$$\liminf_{n\to\infty}\left(1+\frac{n(C_2+2)+C_0/2}{\sum_{i=1}^n\log^+(a_i(x))}\right)^{-1} = 1.$$

Combining this with (3.1) below finishes the proof. For the proof of the equality

$$\dim_{\mathrm{H}}(\mathscr{J}) = 1/2$$

we refer to [Mun12]. □

Proof of Fact 1.3. We prove that if $a_i(x) \leq K$ for all $i \in \mathbb{N}$, then $x \notin \mathscr{F}_1$. By (2.3) we have for $a_i \geq 0$

$$2\log^+(a_i)+C_1 \leq l(\xi_i).$$

Then we have by (2.1)

$$\begin{aligned}\limsup_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{d\left(B_1\cdots B_n\left(i\right),i\right)} &\leq \limsup_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{\sum_{i=1}^{n}l(\xi_i)}\\ &\leq \limsup\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{2\sum_{i=1}^{n}\log^{+}(a_i(x))+nC_1}\\ &\leq \left(1+\frac{C_1}{2\log(K)}\right)^{-1}<1.\end{aligned}$$

□

Proof of Fact 1.4. It is well known that $\{\alpha\in\mathbb{R}\mid \mathscr{F}_\alpha\neq\varnothing\}$ is an interval (cf. [Sch99]). By the Facts 1.2 and 1.1 we have $\mathscr{F}_0\neq\varnothing$ and $\mathscr{F}_1\neq\varnothing$. Further, we clearly have on the one hand $0\leq\liminf_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{d(B_1\cdots B_n(i),i)}$ and on the other hand, using (2.1) and (2.3), we estimate for all $x\in L_r\left(G\right)$,

$$\limsup_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{d\left(B_1\cdots B_n\left(i\right),i\right)}\leq\limsup_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(a_i(x))}{\sum_{i=1}^{n}l(\xi_i)}\leq\lim_{n\to\infty}\frac{\sum_{i=1}^{n}l(\xi_i)}{\sum_{i=1}^{n}l(\xi_i)}=1. \tag{3.1}$$

This proves the fact. □

4. Proof of Theorem 1.5

The radial limit set $L_r(G)$ of a free Fuchsian group G has a representation as an infinite conformal graph directed Markov system ([MU03]). For the details, we refer to [KS04].

Multifractal formalism for the interior of the spectrum. The multifractal formalism can be established as in [JK10] (see also [KS04]). The formula for the Hausdorff dimension for the interior of the spectrum also follows by extending the arguments given in [JK11] for iterated function systems to graph directed Markov systems.

These observations lead to the following result from the multifractal formalism as provided in, for instance, [KS04, KS07, JK10]. Set $\alpha_0:=\left(t'\right)^{-1}(0)$, where $\left(t'\right)^{-1}$ denotes the inverse function of t' and let

$$\mathscr{F}_\alpha^*:=\begin{cases}\left\{x\in L_r\left(G\right)\mid\limsup_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(|B_i|-1)}{d(B_1\cdots B_n(i),i)}\geq\alpha\right\}, & \alpha\geq\alpha_0,\\ \left\{x\in L_r\left(G\right)\mid\liminf_{n\to\infty}\frac{2\sum_{i=1}^{n}\log^{+}(|B_i|-1)}{d(B_1\cdots B_n(i),i)}\leq\alpha\right\}, & \alpha<\alpha_0.\end{cases}$$

Then we have for $\alpha\in(0,1)$

$$f(\alpha)=\dim_{\mathrm{H}}\left(\mathscr{F}_\alpha\right)=\dim_{\mathrm{H}}\left(\mathscr{F}_\alpha^*\right)=-\widehat{t}(-\alpha). \tag{4.1}$$

Here, we used Lemma 2.4 to express the level sets in $\mathscr{F}_\alpha^*\cap\mathscr{D}$ as quotients of Birkhoff sums, and the observation that $\mathscr{F}_\alpha^*$ and $\mathscr{F}_\alpha^*\cap\mathscr{D}$ have the same Hausdorff dimension. We can make this observation because $\mathscr{F}_\alpha^*$ is a countable union of bi-Lipschitz images of $\mathscr{F}_\alpha$.

Asymptotic behaviour for the left boundary point. By Fact 1.1 we have

$$\dim_{\mathrm{H}}\left(\mathscr{F}_0\right)\geq\dim_{\mathrm{H}}\left(L_c\right)=\delta_c.$$

Combining (4.1) and Lemma 2.6 the lower bound follows from

$$\dim_{\mathrm{H}}\left(\mathscr{F}_0\right)\leq\lim_{\alpha\to0}\dim_{\mathrm{H}}\left(\mathscr{F}_\alpha^*\right)=\delta_c.$$

Asymptotic behaviour for the right boundary point. Combining Fact 1.2 and 1.3 we obtain $\dim_{\mathrm{H}}(\mathscr{F}_1) \geq \dim_{\mathrm{H}}(\mathscr{J}) = 1/2$.

Now this observation and the fact that $\mathscr{F}_1 \subset \mathscr{F}_\alpha^*$ for every $\alpha_0 < \alpha < 1$ together with Lemma 2.7 finally gives

$$\dim_{\mathrm{H}} \mathscr{F}_1 \leq \lim_{\alpha \nearrow 1} \dim_{\mathrm{H}} \mathscr{F}_\alpha^* \leq 1/2.$$

This proves the claim for the properties of the right boundary point of the spectrum.

The derivatives in the boundary points. For the derivative of f at the boundary points we use the general identity for Legendre transforms

$$f'(\alpha) = \left(\widehat{t}\right)'(-\alpha) = -\left(t'\right)^{-1}(-\alpha), \ \ \alpha \in (0,1).$$

The real-analyticity then gives

$$\lim_{\alpha \searrow 0} f'(\alpha) = -\lim_{\alpha \nearrow 1} f'(\alpha) = +\infty.$$

Irregular set. The fact that the set of points for which the mean cusp winding number does not exist has full Hausdorff dimension $\delta(G)$ follows from [BS00] by exhaustion of Σ_A with finite alphabet subsystems.

REFERENCES

[BJ97] C. J. Bishop and P. W. Jones, *Hausdorff dimension and Kleinian groups*, Acta Math. **179** (1997), no. 1, 1–39. MR 1484767 (98k:22043)

[BS79] R. Bowen and C. Series, *Markov maps associated with Fuchsian groups*, Inst. Hautes Études Sci. Publ. Math. (1979), no. 50, 153–170. MR 556585 (81b:58026)

[BS00] L. Barreira and J. Schmeling, *Sets of "non-typical" points have full topological entropy and full Hausdorff dimension*, Israel J. Math. **116** (2000), 29–70. MR 1759398

[EP94] D. B. A. Epstein and C. Petronio, *An exposition of Poincaré's polyhedron theorem*, Enseign. Math. (2) **40** (1994), no. 1-2, 113–170. MR 1279064

[JK10] J. Jaerisch and M. Kesseböhmer, *The arithmetic-geometric scaling spectrum for continued fractions*, Ark. Mat. **48** (2010), no. 2, 335–360. MR 2672614 (2011g:11144)

[JK11] ———, *Regularity of multifractal spectra of conformal iterated function systems*, Trans. Amer. Math. Soc. **363** (2011), no. 1, 313–330. MR 2719683

[KS04] M. Kesseböhmer and B. O. Stratmann, *A multifractal formalism for growth rates and applications to geometrically finite Kleinian groups*, Ergodic Theory Dynam. Systems **24** (2004), no. 1, 141–170. MR 2041265 (2005e:37061)

[KS07] ———, *A multifractal analysis for Stern-Brocot intervals, continued fractions and Diophantine growth rates*, J. Reine Angew. Math. **605** (2007), 133–163. MR 2338129 (2008f:37018)

[MU03] R. D. Mauldin and M. Urbański, *Graph directed Markov systems*, Cambridge Tracts in Mathematics, vol. 148, Cambridge University Press, Cambridge, 2003, Geometry and dynamics of limit sets. MR MR2003772 (2006e:37036)

[Mun12] S. Munday, *On Hausdorff dimension and cusp excursions for Fuchsian groups*, Discrete Contin. Dyn. Syst. **32** (2012), no. 7, 2503–2520. MR 2900557

[Sch99] J. Schmeling, *On the completeness of multifractal spectra*, Ergodic Theory Dynam. Systems **19** (1999), no. 6, 1595–1616. MR 1738952

[SS05] M. Stadlbauer and B. O. Stratmann, *Infinite ergodic theory for Kleinian groups*, Ergodic Theory Dynam. Systems **25** (2005), no. 4, 1305–1323. MR 2158407 (2006e:37010)

[Sta04] M. Stadlbauer, *The return sequence of the Bowen-Series map for punctured surfaces*, Fund. Math. **182** (2004), no. 3, 221–240. MR 2098779 (2006c:37031)

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, SHIMANE UNIVERSITY, NISHIKAWATSU 1060, MATSUE, SHIMANE, 690-8504 JAPAN.

E-mail address: jaerisch@riko.shimane-u.ac.jp

URL: www.math.shimane-u.ac.jp/~jaerisch/

FACHBEREICH 3 – MATHEMATIK UND INFORMATIK, UNIVERSITÄT BREMEN, BIBLIOTHEKSTR. 1, 28359 BREMEN, GERMANY

E-mail address: mhk@math.uni-bremen.de

URL: www.math.uni-bremen.de/stochdyn

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, VIA ZAMBONI, 33, 40126 BOLOGNA, ITALY

E-mail address: saraann.munday@unibo.it